



# Coricidin<sup>®</sup> HBP Cold & Flu

## Drug Facts

<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg.....	Antihistamine

### Uses

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - sore throat
  - sneezing
  - runny nose
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use to sedate children

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### When using this product

- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



## Coricidin® HBP Cold & Flu

**Directions**

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 tablets with water every 4-6 hours. Do not exceed 10 tablets in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Inactive ingredients** colloidal silicon dioxide, FD&C red #40 aluminum lake, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, purified water, stearic acid, talc, titanium dioxide

**Questions or comments? 1-800-317-2165** (Mon-Fri 9AM - 5PM EST)