



Coricidin HBP Nighttime Multi-Symptom Cold Liquid

Drug Facts

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

Uses ▪ temporarily relieves common cold or flu symptoms:
▪ headache ▪ sore throat ▪ runny nose ▪ minor aches and pains
▪ cough due to minor throat and bronchial irritation ▪ sneezing ▪ fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 120 mL (4 doses) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening ▪ blisters ▪ rash ▪ hives
- facial swelling ▪ asthma (wheezing) ▪ shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children

Do not use

- with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough with excessive phlegm (mucus)
- glaucoma
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if



Coricidin HBP Nighttime Multi-Symptom Cold Liquid

- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- redness or swelling is present

If **pregnant or breast-feeding**, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than recommended dose
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 6 hours while symptoms last
- do not exceed 120 mL (4 doses) in 24 hours, unless directed by a doctor
- children under 12 years of age: do not use

Other information

- **each 30 mL contains:** sodium 39 mg
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue No.1, FD&C red No.40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-317-2165 (Mon – Fri 9AM – 5PM EST)