



# Coricidin<sup>®</sup> HBP Cough & Cold

<b>Drug Facts</b>	
<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
Chlorpheniramine maleate 4 mg.....	Antihistamine
Dextromethorphan hydrobromide 30 mg.....	Cough suppressant
<b>Uses</b>	
<ul style="list-style-type: none"><li>temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold</li><li>temporarily relieves runny nose and sneezing due to the common cold</li></ul>	
<b>Warnings</b>	
<b>Do not use</b>	
<ul style="list-style-type: none"><li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li><li>if you have ever had an allergic reaction to this product or any of its ingredients</li><li>in children under 12 years of age</li></ul>	
<b>Do not use to sedate children</b>	
<b>Ask a doctor before use if you have</b>	
<ul style="list-style-type: none"><li>glaucoma</li><li>difficulty in urination due to enlargement of the prostate gland</li><li>cough that occurs with excessive phlegm (mucus)</li><li>a breathing problem such as emphysema or chronic bronchitis</li><li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li></ul>	
<b>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers</b>	
<b>When using this product</b>	
<ul style="list-style-type: none"><li>excitability may occur, especially in children</li><li>marked drowsiness may occur</li><li>avoid alcoholic beverages</li><li>alcohol, sedatives and tranquilizers may increase drowsiness</li><li>use caution when driving a motor vehicle or operating machinery</li></ul>	
<b>Stop use and ask a doctor if cough persists for more than 7 days, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.</b>	
<b>If pregnant or breast-feeding, ask a health professional before use.</b>	
<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	
<ul style="list-style-type: none"><li>do not take more than recommended dose</li><li>adults and children 12 years and over: take 1 tablet with water every 6 hours. Do not exceed 4 tablets in 24 hours or as directed by a doctor.</li><li>children under 12 years of age: do not use</li></ul>	
<b>Other information</b>	
<ul style="list-style-type: none"><li>store between 20° to 25°C (68° to 77°F)</li></ul>	
<b>Inactive ingredients</b> croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C yellow no. 6 aluminum lake, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide	
<b>Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)</b>	