



# Coricidin HBP

## Maximum Strength Multi-Symptom Flu

<b>Drug Facts</b>	
<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg.....	Antihistamine
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant

**Uses**

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - cough
  - runny nose
  - sneezing
  - sore throat
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

---

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

---

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with excessive phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

---

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

---

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

---

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur



## Coricidin HBP Maximum Strength Multi-Symptom Flu

<ul style="list-style-type: none"><li>▪ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li></ul> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children. Abuse of this product can lead to serious injury.</b></p> <p><b>Overdose warning:</b> Taking more than the recommended dose may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>
<p><b>Directions</b></p> <ul style="list-style-type: none"><li>▪ do not use more than directed (see overdose warning)</li><li>▪ adults and children 12 years and over: take 2 tablets every 4 hours, while symptoms persist. Do not take more than 10 tablets in 24 hours or as directed by a doctor.</li><li>▪ children under 12 years of age: ask a doctor</li></ul>
<p><b>Other information</b></p> <ul style="list-style-type: none"><li>▪ store at room temperature. Avoid excessive heat.</li></ul>
<p><b>Inactive ingredients</b> FD&amp;C red #40 aluminum lake, lecithin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide</p>
<p><b>Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)</b></p>